

膵 GEM療法

軽 リスク

		1クール 4 週間																												
		day																												
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
ゲムシタピン注 1000mg/m ² (day1,8,15)																														

膵 FOLFIRINOX療法

高 リスク

		1クール 2 週間													
		day													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
オキサリプラチン注 85mg/m ² (day1)															
イリノテカン注 180mg/m ² (day1)															
レボホリナート注 200mg/m ² (day1)															
フルオロウラシル注 400mg/m ² (day1)															
フルオロウラシル注 2400mg/m ² (day1-2)															
アプレピタントCap 125mg	1														
アプレピタントCap 80mg		1	1												
デカドロン錠4mg (2錠/2×MT)		2	2	2											

膵 mFOLFIRINOX療法

高 リスク

		1クール 2 週間													
		day													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
オキサリプラチン注 85mg/m ² (day1)															
イリノテカン注 150mg/m ² (day1)															
レボホリナート注 200mg/m ² (day1)															
フルオロウラシル注 2400mg/m ² (day1-2)															
アプレピタントCap 125mg	1														
アプレピタントCap 80mg		1	1												
デカドロン錠4mg (2錠/2×MT)		2	2	2											

膵 GEM+S-1療法

軽 リスク

		1クール 3 週間																								
		day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21			
S-1 80mg/m ² /day (day1-14)		2週間内服																								
ゲムシタビン注 1000mg/m ² (day1,8)																										

S-1 1日量/body
 <1.25m² → 80mg/day
 1.25-1.5m² → 100mg/day
 ≥1.5m² → 120mg/day

膵 GEM+Nab-PAC療法

軽 リスク

		1クール 4 週間																												
		day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
ゲムシタビン注 1000mg/m ² (day1,8,15)																														
アブラキサン注 125mg/m ² (day1,8,15)																														

膵 nal-IRI+5FU/LV療法

中① リスク

		1クール 2 週間														
		day	1	2	3	4	5	6	7	8	9	10	11	12	13	14
オニバイド注 70mg/m ² (day1)																
レボホリナート注 200mg/m ² (day1)																
フルオロウラシル注 2400mg/m ² (day1-2)																
アプレピタントCap 125mg		1														
アプレピタントCap 80mg			1	1												